

## 症 例 報 告

## 八味地黄丸により軽快した下腹部痛の1症例

榎田康彦\* 大成美華\* 上田智佳子\*  
喜多正樹\* 山本 健\* 小林 勉\*

要旨：慢性疼痛症例では、鎮痛薬への依存を生じる可能性がある。症例は、27歳女性。1996年2月より、左下腹部痛が出現し、左卵巣腫大を認めたため同年3月に部分切除が施行された。それ以外に器質的病因は認められなかったが、疼痛は消失せず、1998年3月に紹介された時には、ペンタゾシン依存が生じていた。八味地黄丸の内服と1～2週間に1回のケタミン静注療法によりペンタゾシン依存から脱却した。

索引用語：慢性疼痛、ペンタゾシン、八味地黄丸、ケタミン、下腹部痛

## PAIN AND KAMPO MEDICINE Vol.9 1999

A case of lower abdominal pain successfully treated with Hachimi-jio-gan  
Yasuhiko Kushida\*, Mika Ohnari\*, Tikako Ueda\*, Masaki Kita\*,  
Ken Yamamoto\* and Tsutomu Kobayashi\*

**Abstracts** : It is possible that patients with chronic pain to be addicted to analgesics. A 27-year-old female had felt left lower abdominal pain since February 1996. She was diagnosed as having swelling of the left ovarium and underwent a partial resection of the left ovarium. There was no abnormalities except the swelling. But her abdominal pain did not disappear after the operation. When she consulted our clinic in March 1998, she had been addicted to pentazocine. We started to give her Hachimi-jio-gan regularly and ketamine once per week or two weeks. which succeeded to cure her addiction.

**Key words** : chronic pain, pentazocine, Hachimi-jio-gan, ketamine, lower abdominal pain

\* Department of Anesthesiology and Critical Care Medicine, School of Medicine, Kanazawa University  
Offprint requests to : Yasuhiro Kushida, Department of Anesthesiology and Critical Care Medicine, School of Medicine, Kanazawa University. 13-1 Takara-machi, Kanazawa 920-8641, Japan