

症 例 報 告

## failed back surgery syndrome に対して 抑肝散加陳皮半夏が有効であった1症例

竹田 貴雄\*<sup>1</sup> 増田 将史\*<sup>1</sup> 平原 志保\*<sup>2</sup>  
西村 昌泰\*<sup>1</sup> 岡野 祐子\*<sup>1</sup> 岡野 龍介\*<sup>1</sup>

**要旨：**症例は86歳女性。既往歴に脳梗塞。腰部脊柱管狭窄症術後も続く腰下肢痛 (failed back surgery syndrome) に対して仙骨硬膜外ブロック，鍼治療を行っていたが，症状に変化はなかった。脳梗塞後も続く顔面と上肢のしびれに対して，たまたま抑肝散加陳皮半夏 (TJ-83) を処方したところ，しびれの改善と共に腰下肢痛も軽快した。

**索引用語：**failed back surgery syndrome, 抑肝散加陳皮半夏

### PAIN AND KAMPO MEDICINE Vol.13 (2003)

#### A case of Failed Back Surgery Syndrome Effectively Treated With Yokukansan-Ka-Chinpi-Hange (TJ-83)

Takao Takeda \*<sup>1</sup>, Masashi Masuda \*<sup>1</sup>, Shiho Hirahara \*<sup>2</sup>,  
Masahiro Nishimura \*<sup>1</sup>, Yuko Okano \*<sup>1</sup>, Ryusuke Okano \*<sup>1</sup>

**Abstract:** An 86 year-old female with a past history of cerebral infarction suffered from postoperative painful symptoms due to lumbar spinal canal stenosis (failed back surgery syndrome). Neither caudal epidural block or acupuncture were effective in relieving her back and leg pain. Three months later, Yokukansan-ka-chinpi-hange (TJ-83) was administered for the treatment of facial and brachial paresthesia persisting after cerebral infarction. With relief of facial and brachial paresthesia, back and leg pain was also relieved unexpectedly after the administration of TJ-83.

**Key words:** failed back surgery syndrome, Yokukansan-ka-chinpi-hange

---

\*<sup>1</sup> Department of Anesthesiology, Nippon Steel Hirohata Hospital

*Offprint requests to:* Takao Takeda, Department of Anesthesiology, Nippon Steel Hirohata Hospital, 3-1, Yumesaki-cho, Hirohata-ku, Himeji-city, Hyogo Pref. 671-1122, Japan

Tel: +81-792-37-6817

\*<sup>2</sup> Department of Anesthesiology, University of Occupational and Environmental Health, Japan