

症 例 報 告

イレウス症状を伴う癌性腹膜炎に対する大建中湯の効果

角田 純一* 水口 公信*

要旨：イレウス症状を伴う癌性腹膜炎に対する大建中湯（TJ-100）の効果を検討した。対象を緩和ケア入院中の、癌性腹膜炎で症状緩和の治療を受けているものとした。終末期癌患者のイレウス症状を伴う癌性腹膜炎は、嘔気・嘔吐、腹痛、腹部膨満などの不快な身体症状を起こすことが多い。今回、そのような症例4例に大建中湯の投与を試み、効果が得られたので報告し考察を加えた。大建中湯は、イレウス症状を伴う癌性腹膜炎に対して投与を試みる方法の一つと考えられた。

索引用語：癌性腹膜炎、イレウス症状、大建中湯

PAIN AND KAMPO MEDICINE Vol.13 (2003)

The effect of Dai-kenchu-to on terminal patients with peritoneal carcinomatosis and ileus

Junichi Tsunoda*, Tadanobu Mizuguchi

Abstract: The effect of Dai-kenchu-to was investigated in terminal patients with peritoneal carcinomatosis and ileus. Subjects were treated with Dai-kenchu-to. Peritoneal carcinomatosis is defined as the spread and implantation of tumor cells through the peritoneal cavity leading to significant suffering. Major complications of peritoneal carcinomatosis include partial and complete obstruction of the gastrointestinal tract, leading to crampy abdominal pain, nausea, vomiting and abdominal distension. Palliative management of peritoneal carcinomatosis and ileus was performed with Dai-kenchu-to in four terminal patients. It is concluded that the effect of Dai-kenchu-to is beneficial in alleviating the symptom of peritoneal carcinomatosis and ileus.

Key words: peritoneal carcinomatosis, ileus, Dai-kenchu-to

* Palliative Care, Sannou Hospital

Offprint requests to: Junichi Tsunoda, Palliative Care, Sannou Hospital 166-2, Sanno-cho, Inage-ku, Chiba-shi, 263-0002, Japan