

症 例 報 告

子宮悪性腫瘍術後の遷延性下腹部痛に対する漢方治療

林 明宗 *

要旨：子宮悪性腫瘍摘出術および術後放射線治療を受けた後に下腹部痛が30年間遷延した患者に対して、当帰四逆加呉茱萸生姜湯加修治附子末Nが奏効した一例を報告した。疼痛の発症原因としては、悪性腫瘍治療後の骨盤内組織の癒着や瘢痕形成に起因する骨盤内静脈叢の血流障害（芎血）により疼痛症状が出現し、加齢にともなってこれが悪化したものと推測された。

索引用語：子宮悪性腫瘍, 術後下腹部痛, 当帰四逆加呉茱萸生姜湯, 修治附子末N, 芎血

PAIN AND KAMPO MEDICINE Vol.15 (2005)

Kampo Therapy for Prolonged Intractable Lower-abdominal Pain after the Radical Therapy of Malignant Uterine Tumor.

Akimune Hayashi *

Abstract: A 66-year-old woman had radical surgery and postoperative radiotherapy for a malignant uterine tumor. She suffered from postoperative intractable lower abdominal pain for 30 years. She was satisfactorily treated with tokishigyakukagoshuyushokyoto with heat processed Aconite Tuber Powder N. Disturbed intrapelvic blood circulation (blood stasis) due to tissue adhesion and scar formation was considered as the causes of her lower abdominal pain. And aging deteriorated this pathological condition.

Key words: malignant uterine tumor, post operative abdominal pain, Tokishigyakukagoshuyushokyoto, Aconite Tuber Powder, blood stasis

* Department of Neurosurgery, Kanagawa Cancer Center. Division of Kanpo Medicine, Department of General Internal Medicine, Yokohama City University Medical Center

Offprint requests to: Akimune Hayashi, Department of Neurosurgery, Kanagawa Cancer Center. Division of Kanpo Medicine, Department of General Internal Medicine, Yokohama City University Medical Center. 1-1-2 Nakao, Asahi-Ku, Yokohama 241-0815, Japan