

症 例 報 告

頸椎捻挫による後頭神経痛への漢方治療 2 例 — 疎経活血湯による著効・桂枝加朮附湯による有効 —

林 俊樹*¹ 林 智樹*¹ 高口眞一郎*^{1,2} 寺岡俊人*²

要旨：頸椎捻挫による後頭神経痛に対する漢方薬（疎経活血湯他）の効果について述べる。

1) 中間証で芎血のある，39歳男性の後頭神経痛にはツムラ疎経活血湯（TJ-53）で，しびれにはTJ-53+ツムラ桂枝加朮附湯（TJ-18）で，著効をみた。

2) 胃内停水と軽度の芎血のある，30歳女性の後頭神経痛には（五苓散他を約3週用いた後），ツムラ桂枝加朮附湯が有効であったが，略治には数回の星状神経ブロックを必要とした。

索引用語：頸椎捻挫による後頭神経痛，疎経活血湯，桂枝湯類

PAIN AND KAMPO MEDICINE Vol.15 (2005)

Kampo Medicines for Occipital Neuralgia caused by Cervical Sprain

— a markedly effective case by Sokeikakketsuto;

a moderately effective case with Keishikajutsu —

Toshiki Hayashi*¹, Tomoki Hayashi*¹, Shinichiro Takaguchi*^{1,2} and Toshito Teraoka*²

Abstract: We address the effect of Kampo medicines (sokeikakketsuto etc.) on occipital neuralgia caused by cervical sprain.

1) Sokeikakketsuto (TJ-53) was administered to a 39-years-old male patient with moderate physical strength and blood stagnation for occipital neuralgia. It was markedly effective. TJ-53 and keishikajutsu (TJ-18) were given to him and they were markedly effective to *sibire* (numbness)

2) Keishikajutsu was administered to a 30-years-old female patient with intragastric stagnation of water and slightly blood stagnation for occipital neuralgia. It was effective after the 3-week medication of goreisan etc. Further, her symptom almost disappeared by several times of stellate ganglion block.

Key words: occipital neuralgia caused by cervical sprain, Sokeikakketsuto, genera of Keishito

*¹ Marugame Daiichi Hospital

Offprint requests to: Shinichiro Takaguchi, Marugame Daiichi Hospital, 91-3, Tor-machi, Marukame-City, Kagawa 763-0043, Japan