## 症例報告

## 加味逍遙散、補中益気湯を併用した乳癌術後創部痛の一症例

## 大木 浩\*

**要旨**: 手術後の創部痛は非ステロイド性消炎鎮痛剤の投与や時間の経過と共に軽快することがほとんどである. 今回, 乳癌手術後に発症した, 7年も持続した突き刺すような性質の創部痛に対してオピオイド系鎮痛薬, 神経ブロック療法の効果が十分でなく, 西洋薬とともに加味逍遙散 (TJ-24), 補中益気湯 (TJ-41) の併用が有効だった症例を経験したので報告する.

索引用語:乳癌術後痛、加味逍遙散、補中益気湯

## PAIN AND KAMPO MEDICINE Vol. 21 (2011)

Combination treatment of kamishoyosan and hochuekkito on the postoperative wound pain in a breast cancer patient Hiroshi Oki\*

**Abstract**: Postoperative wound pain is usually relieved by the treatment with nonsteroidal anti-inflammatory drugs or with the time passage. We experienced a case that the opioid therapy or the nerve block showed inadequate effect on the treatment of a stabbing pain after breast cancer surgery which continued for 7 years and more, but the combination of kamishoyosan (TJ-24) and hochuekkito (TJ-41) was effective.

Key words: Wound pain after breast cancer surgery, kamishoyosan, hochuekkito

<sup>\*</sup> Department of Anesthesiology, Fujisawa Shounandai Hospital, Japan Offprint requests to: Hiroshi Oki, Department of Anesthesiology, Fujisawa Shounandai Hospital. 2345 Takakura Fujisawa-City, 252-0802, Japan.