

症 例 報 告

脊髄終糸症候群手術後の頭痛に五苓散が即効した 1 症例

坪 敏仁^{*1} 橋場英二^{*1} 大川浩文^{*2} 廣田和美^{*2}

要旨：終糸症候群手術後の頭痛に対して五苓散 (TJ-17) を使用し著効を得た。五苓散はフェンタニル 2.5 μ g 以上の鎮痛効果を示した。五苓散 (TJ-17) の効果発現時間は 20 分と短く、単に脳脊髄液圧に対する効果とは考えにくかった。頭痛改善の機序などについて考察を加え報告する。

索引用語：五苓散, 頭痛, 低脳脊髄圧

PAIN AND KAMPO MEDICINE Vol.24 (2014)

Goreisan was effective to the headache after tight filum terminate operation within 20 minutes.

Toshihito TSUBO^{*1}, Eiji HASHIBA^{*1}, Hirobumi OKAWA^{*2}
and Kazumi HIROTA^{*2}

Abstract: Goreisan was effective to the headache after tight filum terminate operation. It began to work within 20 minutes after its administration. We thought its effect appeared too quick and investigated how Goreisan work.

Key words: Goreisan, head ache,

^{*1} Intensive Care Unit, University of Hirosaki, School of Medicine

Offprint requests to: Toshihito TSUBO, Intensive Care Unit, University of Hirosaki, School of Medicine,
53 Honcho, Hirosaki, Aomori 036-8203, Japan

^{*2} Department of Anesthesiology, University of Hirosaki, School of Medicine