

症 例 報 告

胸部解離性大動脈瘤に対する手術を契機に発症した 手指のしびれ，痛みに疎経活血湯が有効であった1症例

渡邊秀和^{*1,*2} 滝口規子^{*1} 千葉知史^{*1} 綿引奈苗^{*1} 北村知子^{*1}
堤 祐介^{*1} 山城 晃^{*1} 伊達 久^{*1} 関 隆志^{*2,*3}

要旨：症例は70歳代男性。7年ほど前の胸部解離性大動脈瘤に対する手術後から左手指にしびれを感じるようになった。しびれは次第に増悪し痛みも感じるようになった。また，その範囲も両側の手指に拡大した。手術を契機に血瘀，血虚となり，血虚に乗じて風寒湿邪が侵入し気血を阻滞することにより，しびれ，痛みが生じたものと考え，疎経活血湯を開始したところしびれ，痛みは軽減した。その後，腎虚も認められたため八味地黄丸や牛車腎気丸を併用したところ症状は消失した。

索引用語：疎経活血湯，牛車腎気丸，風湿痺，右大動脈弓，解離性大動脈瘤

PAIN AND KAMPO MEDICINE Vol.28 (2018)

Successful treatment with Sokeikakketsuto for finger pain and numbness caused by the operation for dissecting aneurysm of the aorta with right aortic arch.

Hidekazu WATANABE^{*1,2}, Noriko TAKIGUCHI^{*1}, Tomofumi CHIBA^{*1},
Nanae WATABIKI^{*1}, Tomoko KITAMURA^{*1}, Yusuke TSUTSUMI^{*1},
Akira YAMASHIRO^{*1}, Hisashi DATE^{*1} and Takashi SEKI^{*2,3}

Abstract: A 70-year-old man was a case. About seven years ago, he came to feel numbness in the left hand finger triggered by the operation for chest dissection aortic aneurysm with right aortic arch. Numbness gradually worsened and he felt pain. In addition, the area of numbness also expanded to both hands.

The surgery caused blood deficiency. Blood deficiency easily caused wind-dampness. Wind-dampness made stasis of qi and blood. It is thought that stasis of qi and blood caused the numbness and pain. Sokeikakketsuto reduced the numbness and pain. Combination use of sokeikakketsuto and hachimijogan or goshajinkigan cured these symptoms.

Key words: sokeikakketsuto, goshajinkigan, wind-dampness, right aortic arch, dissecting aneurysm of the aorta

*¹ Sendai Pain Clinic Center

Offprint requests to: Hidekazu WATANABE, Sendai Pain Clinic Center.
3-14-1 Shindenhigashi, Miyagino-ku, Sendai-shi, Miyagi 981-0039, Japan

*² Wakuyacho National Health Insurance Hospital

*³ Cyclotron and Radioisotope Center (CYRIC), Tohoku University