

短 報

## 治打撲一方が脳脊髄液減少症の後頸部痛に有効だった 1 例

光藤 尚\* 田村直俊\* 溝井令一\* 荒木信夫\* 山元敏正\*

**要旨：**脳脊髄液減少症に伴う後頸部痛に治打撲一方が有効だった症例を経験した。脳脊髄液減少症と類似する低頭蓋内圧性頭痛においては症状として後頸部痛を高頻度に認めるが、治打撲一方は後頸部痛に有効であることが知られており、脳脊髄液減少症に伴う後頸部痛においても有効であることが示唆された。

**索引用語：**脳脊髄液減少症, 後頸部痛, 治打撲一方

### PAIN AND KAMPO MEDICINE Vol.31 (2022)

#### Successful treatment with jidabokuippo for posterior neck pain due to cerebrospinal fluid hypovolemia: a case report

Takashi MITSUFUJI\*, Naotoshi TAMURA\*, Yoshikazu MIZOI\*, Nobuo ARAKI\*, and Toshimasa YAMAMOTO\*

**Abstract:** We experienced a case of posterior neck pain due to cerebrospinal fluid hypovolemia in which jidabokuippo was effective. Back neck pain is frequently observed in intracranial hypotension headache similar to cerebrospinal fluid hypovolemia. Jidabokuippo is known to be effective for posterior neck pain and was suggested to be effective in posterior neck pain associated with cerebrospinal fluid hypovolemia.

**Key words:** CSF hypovolemia, neck pain, jidabokuippou

---

\*Department of Neurology, Saitama Medical University Hospital  
Takashi MITSUFUJI, Department of Neurology, Saitama Medical University Hospital.  
38, Morohongo, Moroyama-machi, Iruma-gun, Saitama, 350-0495, Japan